

# Expedited Partner Therapy: Gonorrhea

## A Guide for Partners

### Why am I getting this medicine?

- Your partner is being treated for gonorrhea. **Gonorrhea** is an STI (sexually transmitted infection) that you can get through oral, anal, or vaginal sex.
- A medical provider has given your partner medicine to give to you to treat you for gonorrhea. You may have gonorrhea even without any symptoms. If you do not have gonorrhea, this medicine won't do anything. This medicine is so you don't develop serious health problems or pass along the infection to others. This is called **Expedited Partner Therapy (EPT)**.
- EPT is an easy, fast, and private way for you to be treated. The medicine is safe and works well when used the right way.

### Why should I take this medicine?

- If not treated, gonorrhea can lead to serious health problems.

### Serious health problems that can happen:

#### In Women:

- **Pelvic Inflammatory Disease** - can cause pain and affects the reproductive organs.
- **Ectopic pregnancy** - a pregnancy in the fallopian tubes.
- **Infertility** - not being able to have children.

#### In Men:

- a painful condition in the tubes attached to the testicles that can in rare cases lead to infertility .

### What is this medicine?

- The Medicine that treats gonorrhea is **Suprax** (cefixime). This is a **antibiotic** pill (medicine that help fight infection).
- This medicine will get rid of gonorrhea, but **will not treat other STIs**.
- You should not take this medicine if you are allergic to **Suprax**, or if you have serious health problems. Talk with a medical provider right away to see what treatment is best for you.
- If you have any signs of a more serious infection, get medical care right away. Signs include stomach or pelvic, or testicle pain, fever, nausea, or throwing up.

### How should I take this medicine?

- Take all of your pills as instructed. It is important to finish all of your pills

If you think you might have another STI, visit a CHN clinic to get tested.

### After taking this medicine, what should I do?

- Do not have sex (vaginal, oral, or anal) for 7 days after you and all your partners have finished treatment. It takes a full week for the medicine to work.
- Take all of your pills. Wait for 7 days after taking your last pill to have sex.
- Do not share medicine for gonorrhea with anyone.
- You should be tested again about 3 months after your treatment , even if your partner was treated.

### Are there any side effects?

- Not usually, but there can be side effects.

Common side effects that can happen:	Serious side effects are rare, but can happen:
<ul style="list-style-type: none"><li>• Diarrhea</li><li>• Nausea</li><li>• Stomach aches</li><li>• Headaches</li><li>• Throwing up</li></ul>	<p>Some people may have allergic reactions to the medicine. These reactions can cause:</p> <ul style="list-style-type: none"><li>• Trouble breathing</li><li>• Tightness in chest</li><li>• Closing of throat</li><li>• Swelling of lips or tongue</li><li>• Hives (itchy bumps on the skin)</li></ul>

If you have any of the serious side effects after taking this medicine, call 911 or go to emergency room right away.